

JENIS-JENIS ORGANISME PENGANGGU TUMBUHAN KARANTINA, UMBI LAPIS, PERILAKUAN DAN
NEGARA ASAL.

I. SERANGGA/INSECTS

[illegible]

[illegible]

www.djpp.depkmham.go.id

[illegible]

III. CENDAWAN/FUNGAL

NO.	JENIS OPTIK	JENIS UMBI LAPIS	PERLAKUAN	NEGARA ASAL
1.	<i>Caracorum dudifae</i> Wolfson, Pseudoperle lutea; leaf spot of onion and garlic, witherlip of onion and garlic, leaf spot garlic, leaf spot; onion	Adipon ascorbinatun (bawang merah, shallot), Allium spongiosum (sawung bawang), Allium cepa, Chinese	1. Pymigasi 1000/20 Mucili : a. 40 g / M ² / 28°C 3-4°C/2 hari, a.u. u b. 30 g / M ² / 28°C 3-4°C/2 hari, petan 2. Rodiani : 70 gray	Asia: India Amerika : Argentina, Jamaica

IV. TUNGAU/ MITES

NO.	JENIS OPTK	JENIS UMBI LAPIS	PERLAKUAN	NEGARA ASAL
2	<i>Petrobia latens</i> (Tetranychidae) Brown Wheat Mite	<i>Albion sativum</i> (Barang potato/ kentel) <i>Albion apsa</i> (Barang Potato/ kentel)	1. Fungas (CH ₂ Br) Murni : a. 40 gr / M ² / 25°C 3°C / 2 jam, atau b. 76 gr / M ² / 25°C 3°C / 2 jam, atau 2. Inedasi 70 gray	Eropa: Austria, Denmark, France, Germany, Italy, Netherland, Portugal, Poland, Federal Rusia, Spain Asia: China, India, Republic of Korea Africa: Egypt America: Canada, USA Oceania: Australia
3	<i>Rhizoglyphus oryzae</i> Bull Mite	<i>Akkens pp</i>	1. Fungas (CH ₂ Br) Murni : a. 40 gr / M ² / 25°C 3°C / 2 jam, atau b. 82 gr / M ² / 25°C 3°C / 2 jam, atau c. 76 gr / M ² / 25°C 3°C / 2 jam, atau 2. radasi 70 gray	Eropa: Former USSR, Hungary, Spain, Asia: Taiwan, Japan, Korea America: USA (Texas) Oceania: New Zealand

MENTERI PERTANIAN
REPUBLIK INDONESIA,

SUSWONO